# Clinical and technical factors in endoscopic skull base surgery associated with reconstructive success

Arash Abiri<sup>1</sup>, Benjamin F. Bitner<sup>1</sup>, Theodore V. Nguyen<sup>1</sup>, Jonathan C. Pang<sup>1</sup>, Kelsey M. Roman<sup>1</sup>, Milind Vasudev<sup>1</sup>, Dean D. Chung<sup>1</sup>, Siddhant H. Tripathi<sup>3</sup>, Jacob C. Harris<sup>4</sup>, Nikitha Kosaraju<sup>5</sup>, Ryan M. Shih<sup>6</sup>, Myungjun Ko<sup>6</sup>, Jessa E. Miller<sup>5</sup>, Jennifer E. Douglas<sup>4</sup>, Daniel J. Lee<sup>7</sup>, Jacob G. Eide<sup>8</sup>, Rijul S. Kshirsagar<sup>9</sup>, Katie M. Phillips<sup>3</sup>, Ahmad R. Sedaghat<sup>3</sup>, Marvin Bergsneider<sup>6</sup>, Marilene B. Wang<sup>5</sup>, James N. Palmer<sup>4</sup>, Nithin D. Adappa<sup>4</sup>, Frank P.K. Hsu<sup>2</sup>, Edward C. Kuan<sup>1,2</sup>

Rhinology 62: 3, 330 - 341, 2024 https://doi.org/10.4193/Rhin23.267

# Abstract

**Background**: In this study, we identified key discrete clinical and technical factors that may correlate with primary reconstructive success in endoscopic skull base surgery (ESBS). **Methods**: ESBS cases with intraoperative cerebrospinal fluid (CSF) leaks at four tertiary academic rhinology programs were retrospectively reviewed. Logistic regression identified factors associated with surgical outcomes by defect subsite (anterior cranial fossa [ACF], suprasellar [SS], purely sellar, posterior cranial fossa [PCF]). **Results**: Of 706 patients (50.4% female), 61.9% had pituitary adenomas, 73.4% had sellar or SS defects, and 20.5% had high-flow intraoperative CSF leaks. The postoperative CSF leak rate was 7.8%. Larger defect size predicted ACF postoperative leaks; use of rigid reconstruction and older age protected against sellar postoperative leaks; and use of dural sealants compared to fibrin glue protected against PCF postoperative leaks. SS postoperative leaks occurred less frequently with the use of dural onlay. Body-mass index, intraoperative CSF leak flow rate, and the use of lumbar drain were not significantly associated with postoperative CSF leak. Meningitis was associated with larger tumors in ACF defects, nondissolvable nasal packing in SS defects, and high-flow intraoperative leaks in PCF defects. Sinus infections were more common in sellar defects with synthetic grafts and nondissolvable nasal packing. **Conclusions**: Depending on defect subsite, reconstructive success following ESBS may be influenced by factors, such as age, defect size, and the use of rigid reconstruction, dural onlay, and tissue sealants.

Key words: endoscopic skull base surgery, skull base reconstruction, skull base repair, risk factors, CSF leak

# **Graphical abstract**

# Endoscopic Skull Base Surgery cases with intraoperative Cerebrospinal Fluid (CSF) leaks Retrospective review at four tertiary academic rhinology programs



Abiri A, Bitner BF, Nguyen TV, et al. Rhinology 2024. https://doi.org/10.4193/Rhin23.267

# Introduction

Over the past two decades, advancements in endoscopic skull base surgery (ESBS) have made it possible to manage many skull base pathologies via a minimally invasive approach that otherwise previously required open resection <sup>(1)</sup>. Furthermore, for properly selected patients, ESBS is associated with decreased surgical morbidity, shorter hospital stays, and faster recovery times, which has facilitated the adoption of endoscopic approaches among otolaryngologists and neurosurgeons at highvolume skull base surgery centers <sup>(2-4)</sup>. However, the incidence of postoperative cerebrospinal fluid (CSF) leak following reconstruction presents a major technical challenge and serves as the primary cause of morbidity following ESBS <sup>(5,6)</sup>.

Various factors, such as defect site, defect size, and intraoperative CSF leak flow rate have been found to predict reconstructive outcomes <sup>(1)</sup>. Numerous surgical techniques and materials have been proposed over the years to improve the likelihood of successful skull base repair and reduce postoperative complications <sup>(1,7,8)</sup>. However, systematic reviews examining ESBS techniques have demonstrated a high degree of heterogeneity across studies, with limited high-quality comparative evidence to suggest an optimal repair technique <sup>(9-11)</sup>. Additionally, there is a paucity of studies investigating the influence of specific reconstruction layers, materials, and adjunctive measures on reconstructive outcomes following ESBS.

In this study, we present one of the largest series to date of endonasal intradural skull base defect repairs. We aimed to identify patient-specific factors (e.g., sex, body mass index, defect site and size), technical factors (e.g., reconstruction layers and materials), and perioperative interventions and adjunctive measures (e.g., lumbar drain, nasal packing) that predict major reconstructive outcomes, including postoperative CSF leak, meningitis, and sinus infection.

#### **Materials and methods**

#### **Study population**

A retrospective chart review of ESBS cases spanning from December 7, 2007 to December 21, 2022 was conducted at four tertiary academic rhinology programs (University of California, Irvine, Orange, CA, USA; University of California, Los Angeles, Los Angeles, CA, USA; University of Pennsylvania, Philadelphia, PA, USA; University of Cincinnati, Cincinnati, OH, USA) in accordance with each institution's corresponding Institutional Review Board. Inclusion criteria included any adult patient (age  $\geq$ 18 years) who underwent ESBS and exhibited an intraoperative CSF leak (Table 1).

#### **Study variables**

Independent covariates used for analysis included overall patient demographics, intraoperative CSF leak flow rate (i.e., low-flow vs. high-flow), defect site, lumbar drain (LD) placement,



Figure 1. Boundaries of skull base defect sites analyzed in the current study: anterior cranial fossa (red), suprasellar (orange), sellar (green), and posterior cranial fossa (blue). The suprasellar (\*) and prepontine (\*\*) cisterns are indicated.

nasal packing use (none/dissolvable vs. nondissolvable), and technical factors, including use of rigid reconstruction, multilayer inlay grafts (subdural/epidural placement), dural onlay (none vs. autologous vs. synthetic), mucosal coverage (none vs. free mucosal graft [FMG] vs. vascularized flap, including nasoseptal flap [NSF]), and tissue sealant use (none vs. fibrin glue vs. dural sealant). CSF leaks were classified based on the scale described by Esposito et al., with low-flow leaks as grade 1 or 2 and highflow leaks as grade 3 (defect with direct extension to suprasellar/ prepontine cistern and/or dural defect > 1 x 1 cm)  $^{(12)}$ . Figure 1 provides definitions of the skull base defect sites as it pertains to this study. Anterior cranial fossa (ACF) defects were defined as those involving the posterior table of the frontal sinus and/or ethmoid roof (e.g., fovea ethmoidalis, lateral lamella, cribriform plate). Suprasellar (SS) defects were defined as those involving the planum sphenoidale and/or tuberculum sella with continuity with the suprasellar cistern, or sellar pathologies requiring removal of bone in those areas. Purely sellar defects only involved the sellar floor. Posterior cranial fossa (PCF) defects were defined as those involving the clivus or craniocervical junction with continuity with the prepontine cistern. For pathologies which overlapped multiple defect subsites, the subsite with highest CSF flow rate and/or largest defect was used (e.g., invasive pituitary adenoma with inferior clival extension was classified as PCF). Primary measured outcomes were postoperative CSF leaks, meningitis, and sinonasal infections (defined as endoscopic evidence of mucopurulence with positive culture with possible treatment with culture-directed antibiotics within 3 months postoperatively).

# **Statistical analysis**

Statistical analyses were performed using R (version 4.2.2; The

Table 1. Summary of demographics, reconstruction technical factors, and postoperative measures and complications (N=706).

Demographics	No. (%)	Reconstruction Factors	No. (%)	Postoperative Measures	No. (%)	Postoperative Complications	<b>No.</b> (%)
Age, yr (mean ± SD)	51.2 ± 17.1	Rigid Reconstruction		Nasal Packing		CSF Leak	
Sex		No	499 (70.7)	None/ Dissolvable	520 (73.9)	No	651 (92.2)
Male	350 (49.6)	Yes	207 (29.3)	Non- dissolvable	184 (26.1)	Yes	55 (7.8)
Female	356 (50.4)	Multi-Layer Inlay		Lumbar Drain		Meningitis	
BMI, kg/m² (mean ± SD)	$29.9 \pm 7.4$	No	504 (71.4)	No	482 (70.5)	No	687 (97.3)
Tumor Size, cm (mean $\pm$ SD)	$2.36 \pm 1.33$	Yes	202 (28.6)	Yes	202 (29.5)	Yes	19 (2.7)
Defect Site		Dural Onlay		Bedrest Days	$2.8 \pm 4.3$	<b>Sinus Infection</b>	
ACF	58 (9.6)	None	514 (79.4)	POD of Debridement	17.6 ± 16.0	No	674 (95.6)
Sella	262 (43.4)	Autologous	14 (2.2)			Yes	31 (4.4)
Suprasellar	181 (30.0)	Synthetic	119 (18.4)				
PCF	103 (17.1)	Mucosal Coverage					
Defect Size, cm <sup>2</sup> (mean ± SD)	$3.27 \pm 2.24$	None	49 (7.0)				
ACF	$3.81 \pm 3.46$	FMG	77 (11.0)				
Sella	2.96 ± 1.71	Vascularized Flap	576 (82.1)				
Suprasellar	3.11 ± 2.08	All Synthetic Grafts					
PCF	4.29 ± 2.17	No	505 (81.3)				
Intraoperative CSF Leak Flow		Yes	116 (18.7)				
Low	561 (79.5)	Tissue Sealant					
High	145 (20.5)	None	279 (40.8)				
Pathologies		Fibrin Glue	72 (10.5)				
Pituitary Adenoma	437 (61.9)	Dural Sealant	291 (42.6)				
Craniopharyngioma	56 (7.9)	Fibrin Glue + Dural Sealant	41 (6.0)				
Meningioma	52 (7.4)						
Rathke's Cleft Cyst	41 (5.8)						
ENB	17 (2.4)						
Chordoma	15 (2.1)						
Encephalocele	15 (2.1)						
Cushing Disease	8 (1.1)						
Other*	65 (9.2)						

ACF: Anterior Cranial Fossa; BMI: Body Mass Index; CSF: Cerebrospinal Fluid; ENB: Esthesioneuroblastoma; FMG: Free Mucosal Graft; POD: Postoperative Day; PCF: Posterior Cranial Fossa; Postoperative Day; SD: Standard Deviation. \* Pathologies representing less than 0.5% of the study population.

R Foundation for Statistical Computing) in RStudio (version 2022.12.0). A p-value of <0.05 was considered statistically significant. Wilcoxon rank sum and chi-square tests were conducted for continuous and categorical variables, respectively, to assess differences in demographics and outcomes between patients with different skull base defects. Logistic regression was conducted to identify predictors of postoperative CSF leak, meningitis, and sinus infection. For analysis of the entire cohort, variables with p-values of <0.1 on univariate logistic regression or those considered clinically significant (e.g., intraoperative leak flow, multilayer inlay, mucosal coverage) based on a priori causal knowledge were included as covariates in multivariable models <sup>(13)</sup>. Models were checked for multicollinearity by ensuring that covariates had variance inflation factors less than 10 <sup>(14)</sup>. For site-specific analysis, due to limits in sample size, only univariate logistic regression was performed.

Variables	Anteri	or Cranial Fos	sa	l	Sellar		S	ıprasellar		Posterio	or Cranial Fos	sa
	No Leak (N = 53)	Leak (N = 5)	٩	No Leak (N = 238)	Leak (N = 24)	٩	No Leak (N = 166)	Leak (N = 15)	۵.	No Leak (N = 94)	Leak (N = 9)	٩
Age, yr	<b>56.0 ± 15.4</b>	46.2 ± 20.8	0.360	50.7 ± 17.7	41.5 ± 15.7	0.013	52.8±15.5	53.2 ± 19.1	0.859	52.6 ± 17.1	49.7 ± 10.7	0.344
Sex, No. (%)												
Male	33 (62.3)	4 (80.0)	0.644	121 (50.8)	10 (41.7)	0.521	78 (47.0)	6 (40.0)	0.788	45 (47.9)	4 (44.4)	>0.99
Female	20 (37.7)	1 (20.0)		117 (49.2)	14 (58.3)		88 (53.0)	(0.0) 9		49 (52.1)	5 (55.6)	
BMI, kg/m²	30.1 ± 6.8	24.3 ± 0.2	0.128	31.6 ± 8.8	35.2 ± 7.3	0.178	28.0 ± 5.6	31.9 ± 7.9	0.174	29.6 ± 7.0	38.1 ± NA	0.198
Tumor Size, cm	3.78 ± 2.07	$4.32 \pm 2.02$	0.481	1.78 ± 1.12	1.88 ± 1.33	0.831	2.42 ± 0.97	2.83 ± 1.44	0.180	2.96 ± 1.28	$2.95 \pm 0.84$	0.756
Defect Size, cm <sup>2</sup>	3.35 ± 2.65	7.71 ± 6.75	0.095	2.88 ± 1.72	3.68 ± 1.56	0.044	3.01 ± 2.02	<b>4.03</b> ± 2.49	0.101	4.22 ± 2.18	4.87 ± 2.18	0.450
Intraop Leak Flow, No. (%)												
Low	30 (56.6)	4 (80.0)	0.392	213 (89.5)	23 (95.8)	0.485	106 (63.9)	(0.0) 9	0.784	76 (80.9)	6 (66.7)	0.384
High	23 (43.4)	1 (20.0)		25 (10.5)	1 (4.2)		60 (36.1)	6 (40.0)		18 (19.1)	3 (33.3)	
Rigid Reconstruction, No. (%)												
No	50 (94.3)	5 (100.0)	>0.99	166 (69.7)	22 (91.7)	0.029	123 (74.1)	14 (93.3)	0.122	69 (73.4)	7 (77.8)	>0.99
Yes	3 (5.7)	0 (0.0)		72 (30.3)	2 (8.3)		43 (25.9)	1 (6.7)		25 (26.6)	2 (22.2)	
Multi-Layer Inlay, No. (%)												
No	32 (60.4)	5 (100.0)	0.148	198 (83.2)	23 (95.8)	0.141	99 (59.6)	11 (73.3)	0.411	72 (76.6)	7 (77.8)	>0.99
Yes	21 (39.6)	0 (0.0)		40 (16.8)	1 (4.2)		67 (40.4)	4 (26.7)		22 (23.4)	2 (22.2)	
Dural Onlay, No. (%)												
None	37 (72.5)	3 (60.0)	0.370	202 (84.9)	22 (91.7)	0.815	104 (67.1)	14 (93.3)	0.008	67 (75.3)	6 (75.0)	0.285
Autologous	3 (5.9)	1 (20.0)		3 (1.3)	0 (0.0)		3 (1.9)	1 (6.7)		2 (2.2)	1 (12.5)	
Synthetic	11 (21.6)	1 (20.0)		33 (13.9)	2 (8.3)		48 (31.0)	0 (0.0)		20 (22.5)	1 (12.5)	
Mucosal Coverage, No. (%)												
None	13 (24.5)	2 (40.0)	0.711	23 (9.7)	2 (8.3)	0.337	0 (0.0)	0 (0.0)	>0.99	0 (0.0)	0 (0.0)	0.576
FMG	16 (30.2)	1 (20.0)		18 (7.6)	4 (16.7)		11 (6.6)	1 (6.7)		8 (8.5)	1 (11.1)	
Vascularized Flap	24 (45.3)	2 (40.0)		197 (82.8)	18 (75.0)		155 (93.4)	14 (93.3)		86 (91.5)	8 (88.9)	
Synthetic Grafts, No. (%)												
No	29 (60.4)	5 (100.0)	0.147	185 (80.8)	24 (100.0)	0.011	96 (69.1)	12 (85.7)	0.235	79 (90.8)	9 (100.0)	>0.99

Table 2. Demographics and reconstruction technical factors in patients with and without postoperative CSF leaks, stratified by defect site.

BMI: Body Mass Index; FMG: Free Mucosal Graft. Percentages reflect the number of patients with available data.

Bold indicates statistically significant, p<0.05

#### Table 2. continued

Variables	Anteri	or Cranial Fos	ia		Sellar		Su	ıprasellar		Posteric	or Cranial Fo	ssa
	No Leak (N = 53)	Leak (N = 5)	٩	No Leak (N = 238)	Leak (N = 24)	٩	No Leak (N = 166)	Leak (N = 15)	۵.	No Leak (N = 94)	Leak (N = 9)	٩
Yes	19 (39.6)	0 (0.0)		44 (19.2)	0 (0.0)		43 (30.9)	2 (14.3)		8 (9.2)	0 (0.0)	
Tissue Sealant, No. (%)												
None	5 (10.6)	2 (40.0)	0.084	82 (38.1)	10 (43.5)	0.882	74 (50.3)	5 (33.3)	0.044	48 (53.9)	4 (50.0)	0.005
Fibrin Glue	11 (23.4)	2 (40.0)		21 (9.8)	2 (8.7)		11 (7.5)	4 (26.7)		5 (5.6)	3 (37.5)	
Dural Sealant	31 (66.0)	1 (20.0)		112 (52.1)	11 (47.8)		62 (42.2)	6 (40.0)		36 (40.4)	1 (12.5)	
Nasal Packing, No. (%)												
None/Dissolvable	22 (41.5)	3 (60.0)	0.643	172 (72.3)	21 (87.5)	0.144	119 (71.7)	11 (73.3)	>0.99	70 (74.5)	9 (100.0)	0.112
Nondissolvable	31 (58.5)	2 (40.0)		66 (27.7)	3 (12.5)		47 (28.3)	4 (26.7)		24 (25.5)	0 (0.0)	
Lumbar Drain, No. (%)												
No	24 (49.0)	2 (40.0)	>0.99	155 (67.7)	16 (66.7)	>0.99	106 (65.8)	8 (57.1)	0.564	71 (77.2)	6 (75.0)	>0.99
Yes	25 (51.0)	3 (60.0)		74 (32.3)	8 (33.3)		55 (34.2)	6 (42.9)		21 (22.8)	2 (25.0)	

# Results

A total of 706 ESBS (50.4% female) patients were studied. The mean age was  $51.2 \pm 17.1$  years. Table 1 lists the sociodemographic and clinical characteristics of this cohort. Overall, 61.9% had pituitary adenomas, 73.4% had sellar or SS defects, and 20.5% had high-flow intraoperative CSF leaks. The average defect size was  $3.27 \pm 2.24$  cm<sup>2</sup>. PCF defects were larger than sella and suprasellar defects (all p<0.05). Postoperatively, 7.8% had CSF leaks, 2.7% had meningitis, and 4.4% had sinus infections. Of the 55 patients with postoperative CSF leaks, 80% had initially presented with low-flow intraoperative leaks and 20% with high-flow leaks.

There were no significant differences in rates of postoperative CSF leak, meningitis, or sinus infection between defect sites (all p>0.05). Defect site-specific differences in demographic and technical reconstruction factors between patients with and without postoperative CSF leak are listed in Table 2. Among sellar defects, larger defects and repairs not involving rigid reconstruction or synthetic grafts more often exhibited postoperative CSF leaks (all p<0.05). The use of dural onlay in SS defect repair and the use of tissue sealants in SS and PCF defect repairs exhibited fewer postoperative leaks (all p<0.05).

Multivariable logistic regression on the entire cohort was used to identify risk factors for postoperative CSF leaks (Table 3). Increased defect size predicted postoperative leak (OR 1.26; 95% CI, 1.04-1.57; p=0.028). Use of rigid reconstruction (OR 0.17; 95% CI, 0.04-0.59; p=0.011), synthetic grafts (OR 0.10; 95% CI, 0.01-0.72; p=0.047), dural sealant (OR 0.14; 95% CI, 0.03-0.50; p=0.005) , and nondissolvable nasal packing (OR 0.22; 95% CI, 0.07-0.60; p=0.005) was protective of postoperative leaks. Sub-analyses stratified by defect site demonstrated that BMI and intraoperative CSF leak flow rate as well as the use of nasal packing, fibrin glue, and lumbar drain were not significantly associated with postoperative CSF leak (all p>0.05).

#### Anterior cranial fossa defects

Of 58 patients with ACF defects, 8.6% had postoperative CSF leak, 5.2% meningitis, and 5.2% sinus infection. On logistic regression, larger defect size was associated with increased risk for postoperative CSF leaks (OR 1.32; 95% Cl: 1.04-1.89; p=0.049). Larger tumors were associated with increased risk for meningitis (OR 1.94; 95% Cl: 1.14-4.06; p=0.028). There were no significant predictors of sinus infection in ACF defects (all p>0.05).

#### **Purely sellar defects**

Of 262 patients with sellar defects, 9.2% had postoperative CSF leak, 2.3% meningitis, and 3.8% sinus infection. Older age (OR 0.97; 95% CI: 0.95-0.99; p=0.017) and rigid reconstruction (OR 0.21; 95% CI: 0.03-0.74; p=0.038) were associated with decreased risk for postoperative CSF leaks. There were no significant predictors of meningitis in sellar defects (all p>0.05). Use of

Table 3. Logistic regression identifying predictors of postoperative CSF leak.

Variables	Univar	iate	Multivari	able
	OR (95% CI)	P-Value	OR (95% CI)	P-Value
Age, yr	0.98 (0.97-0.99)	0.034	0.98 (0.95-1.01)	0.126
Sex				
Male	1 [Reference]		1 [Reference]	
Female	1.20 (0.69-2.09)	0.525	~	~
BMI, kg/m²	1.04 (0.98-1.09)	0.189	~	~
Tumor Size, cm	1.10 (0.89-1.33)	0.370	~	~
Defect Size, cm <sup>2</sup>	1.22 (1.08-1.39)	0.001	1.26 (1.04-1.57)	0.028
Defect Site				
ACF	1 [Reference]		1 [Reference]	
Sella	1.07 (0.42-3.29)	0.897	0.64 (0.13-3.46)	0.590
SS	0.96 (0.35-3.06)	0.936	0.83 (0.15-3.92)	0.836
PCF	1.02 (0.33-3.45)	0.980	0.24 (0.03-1.65)	0.145
Intraoperative CSF Leak Flow Rate				
Low	1 [Reference]		1 [Reference]	
High	0.97 (0.46-1.85)	0.918	1.46 (0.41-4.91)	0.542
Rigid Reconstruction				
No	1 [Reference]		1 [Reference]	
Yes	0.27 (0.10-0.60)	0.003	0.17 (0.04-0.59)	0.011
Multi-Layer Inlay				
No	1 [Reference]		1 [Reference]	
Yes	0.40 (0.17-0.82)	0.020	0.60 (0.12-2.44)	0.501
Dural Onlay				
None	1 [Reference]		1 [Reference]	
Autologous	2.71 (0.60-9.05)	0.136	1.14 (0.11-8.70)	0.90
Synthetic	0.35 (0.10-0.87)	0.045	0.40 (0.03-2.63)	0.41
Mucosal Coverage				
No Flap	1 [Reference]		1 [Reference]	
Vascularized Flap	0.87 (0.45-1.81)	0.680	1.09 (0.25-5.33)	0.909
All Synthetic Grafts				
No	1 [Reference]		1 [Reference]	
Yes	0.15 (0.03-0.50)	0.010	0.10 (0.01-0.72)	0.047
Tissue Sealant				
None	1 [Reference]		1 [Reference]	
Fibrin Glue	2.34 (1.07-4.92)	0.028	0.45 (0.11-1.77)	0.266
Dural Sealant	0.82 (0.43-1.54)	0.532	0.14 (0.03-0.50)	0.005
Nasal Packing				
None/Dissolvable	1 [Reference]		1 [Reference]	
Nondissolvable	0.53 (0.24-1.06)	0.090	0.22 (0.07-0.60)	0.005
Lumbar Drain				
No	1 [Reference]		1 [Reference]	
Yes	1.495 (0.82-2.65)	0.175	1.86 (0.53-6.95)	0.342

ACF: Anterior Cranial Fossa; BMI: Body Mass Index; CI: Confidence Interval; CSF: Cerebrospinal Fluid; OR: Odds Ratio; PCF: Posterior Cranial Fossa; SS: Suprasellar. Bold indicates statistically significant, p<0.05

synthetic grafts (OR 8.15; 95% Cl: 1.79-57.28; p=0.012) and nondissolvable nasal packing (OR 8.53; 95% Cl: 1.89-59.70; p=0.010) were associated with increased risk for sinus infection.

#### Suprasellar defects

Of 181 patients with SS defects, 8.3% had postoperative CSF leak, 1.7% meningitis, and 4.4% sinus infection. The rate of postoperative CSF leak was significantly lower in reconstructions that used a dural onlay (1.9% vs. 11.9%, p=0.035). Use of nondissolvable nasal packing was associated with increased risk for meningitis (OR 5.88; 95% CI: 1.12-43.11; p=0.044). There were no significant predictors of sinus infection in SS defects (all p>0.05).

# Posterior cranial fossa defects

Of 103 patients with PCF defects, 8.7% had postoperative CSF leak, 4.9% meningitis, and 2.0% sinus infection. Fifteen (14.6%) patients presented with chordomas, 5 of whom had high-flow intraoperative leaks, 3 had postoperative CSF leaks, and 2 had meningitis. Compared to fibrin glue, use of dural sealant was associated with reduced risk for postoperative CSF leaks (OR 0.05; 95% Cl: 0.01-0.44; p=0.014). High-flow intraoperative CSF leak was associated with increased risk for meningitis (OR 19.06; 95% Cl: 2.62-385.61; p=0.010). There were no significant predictors of sinus infection in PCF defects (all p>0.05).

#### Discussion

Postoperative CSF leak remains one of the most challenging complications encountered following ESBS. Many reconstructive surgical techniques have therefore been introduced to reduce the incidence of CSF leak, albeit, with variable results. This multicenter study sought to leverage the large number of patients and use of different techniques at four tertiary care rhinology/ anterior skull base surgery programs to investigate patient and technical factors that may be associated with reconstructive success in ESBS. Overall, the incidence of postoperative CSF leak regardless of reconstructive technique used or location of skull base defect was comparable to the current literature, with an overall rate of 7.8%, of which 80% occurred in patients with intraoperative low-flow CSF leaks. This suggests overall highly favorable outcomes with modern techniques and repair strategies. Additionally, there was no difference in CSF leak rate between ACF (8.6%), sella (9.2%), SS (8.3%), or PCF (8.7%) defect locations. Multivariable logistic regression that accounted for potential confounders, such as differences in defect size between subsites, similarly supported a lack of association between defect subsite and CSF leak incidence. Soudry et al.'s systematic review evaluating repair techniques of 673 endoscopically created skull base defects similarly showed that the overall postoperative CSF leak rate after intraoperative skull base repair was 8.5% (57/673)<sup>(15)</sup>. Moreover, this review also demonstrated the location of the skull base defect does not

significantly affect the risk of postoperative CSF leak, excluding clival defects (though the sample size was small).

# Anterior cranial fossa defects

When evaluating factors associated with surgical outcomes by defect subsite, larger ACF defect size was associated with increased risk for postoperative CSF leak. ACF defects can be as large as extending from the posterior table of the frontal sinus to the planum sphenoidale, requiring significant area for coverage, thus increasing the risk for postoperative CSF leaks (16-21). These defects often also require a multilayer reconstruction technique which may include the use of a NSF. With a larger area requiring reconstruction in an anti-gravity configuration, it is conceivable that the forces created by intracranial pressure (ICP) may need to be countered by stiffer materials or bolstering (e.g., nasal packing). Our study does not show any association with onlay used (e.g., NSF vs. FMG) when reconstructing the ACF. The current literature shows variable success with ACF reconstruction; although, recently, rates of postoperative CSF leak are much more consistent and generally low regardless of material used (19,20,22-24). Germani et al. used a nonvascularized single layer to reconstruct the skull base defect with a 3% postoperative CSF leak rate while Eloy et al. reconstructed the ACF with multiple layers including a vascularized flap with a postoperative CSF leak rate of 0% (22,23). Given the variations in reconstructive technique with comparable postoperative outcomes, the success of ACF reconstruction may, in fact, rely on the ability to cover the entire defect regardless of the reconstructive technique used.

#### Sellar defects

There are a myriad of reconstruction options for repairing sellar defects, the most common of ESBS defects, most of which include multiple layers; however, there is no consensus on the optimal reconstructive technique or material (25-27). Some propose graded reconstruction based on the presence of an intraoperative leak and leak flow rate (e.g., high- vs. low-flow) (25,27-35). Most intraoperative CSF leaks following resection of sellar pathologies are low-flow, as demonstrated in this study, with a rate of 90.1% compared to high-flow leaks. In the current study, rigid reconstruction was statistically associated with decreased risk for postoperative CSF leak comprising a rate of 2.7% when used to reconstruct sellar defects compared to a rate of 11.7% when no rigid reconstruction was used. The "gasket seal" technique is one method of rigid reconstruction used for such defects (36,37). It typically consists of a soft overlay (e.g., fascia lata) with a rigid buttress (e.g., bone, synthetic plate) placed into the bony defect and over the overlay creating a gasket that accommodates defect irregularities, thereby creating a watertight seal <sup>(36,37)</sup>. Garcia-Navarro et al. demonstrated success in a larger cohort of 46 patients with a postoperative leak rate of 4.3% (38). Excluding use of rigid reconstruction, no other materials or techniques

were associated with reducing CSF leak. This suggests surgical success is largely independent of specific materials or technique used as long as meticulous technique is employed. Suprasellar defects

Reconstruction of SS defects poses a unique set of challenges. SS defects are typically associated with intraoperative high-flow CSF leaks with continuity with the suprasellar cistern and, in some cases, the third ventricle (39). Difficulty repairing these highflow CSF leaks is amplified when the defect involves multiple skull base planes including the planum sphenoidale, tuberculum sella, and/or sella turcica. Additionally, as the SS defect extends to its limits, the defect border may abut vital structures including the internal carotid arteries and the optic nerves, partially limiting the ability to place inlays <sup>(15)</sup>. In our study, the single most important SS reconstructive technique was the use of a dural onlay, which was associated with a significantly lower rate of postoperative CSF leak. We hypothesize that a dural onlay, which may take the form of autologous (e.g., fascia lata) or synthetic (e.g., porcine small intestine submucosa) graft, may provide an additional layer of watertight coverage over the SS defect, which is commonly irregularly shaped due to exposure of the optic canals and carotid arteries.

# Posterior cranial fossa defects

PCF defects pose unique surgical challenges compared to other skull base defects due to the vertical orientation, limited bony or dural edges, as well as the anatomic proximity to the anterior brain cisterns and ventricles. Reconstruction often requires the use of a vascularized flap for mucosal coverage <sup>(8,40–42)</sup>. Multiple studies have shown clival defects to be particularly difficult to reconstruct; in fact, it has been demonstrated to be the only skull base subsite with improved outcomes using a vascularized flap compared to nonvascularized repair (15,43). One study demonstrated a 60% success rate with multilayer free graft alone whereas the use of a pedicled flap improved success rate to 100% <sup>(44)</sup>. Saito et al. used a fascia lata inlay and NSF onlay with a 100% success rate <sup>(45)</sup>. Two series evaluating resection of clival chordomas with PCF reconstruction with a Duragen inlay and a NSF onlay showed a 90% (9/10) and 75% (15/20) success rate, although the second series comprised more complex patients who were either received prior radiation, had recurrence of disease, or significant intradural extension <sup>(16,46)</sup>. A vascularized flap was used for reconstruction in 91.2% (94/103) of our PCF defects; however, there was no difference in rate of postoperative leak and whether a FMG or vascularized flap was used with leak rates of 11.1% (1/9) and 8.5% (8/94), respectively.

Support materials and adjuncts for reconstruction The benefit of ancillary reconstructive techniques including the use of nasal packing, tissue sealants, and lumbar drains (LD) have long been debated with conflicting outcomes in the literature (47-79). Our data demonstrates a decrease in the risk of CSF leak with the use of dural sealants and nondissolvable nasal packing. Eloy et al. analyzed different tissue sealants including Duraseal® and Tisseel® when applied over a NSF in patients undergoing endoscopic skull base repair for high-flow CSF leaks and compared the incidence of postoperative CSF leak to patients without tissue sealant use (63). This study found there to be no significant difference in the incidence of CSF leak between the two groups. A comprehensive meta-analysis by Ahmed et al. evaluated perioperative LD use following endonasal endoscopic CSF leak repair and found that LD use did not significantly lower postoperative CSF leak recurrence rates (48). There may, however, be certain circumstances in which a LD may be beneficial, most notably high-flow CSF leak in the setting of large ACF and PCF defects (58). Lastly, nasal packing is often used to bolster the reconstruction while the surgical wound heals. The value of packing has been questioned recently with the use of sound techniques and other adjuncts (59,82). Given each of these factors as well as flow rate or other general repair technique did not significantly impact the rate of postoperative CSF leak, success of skull base repair may hinge largely on surgeon adherence to basic principles and necessary elements of skull base reconstruction <sup>(83)</sup>. This is further emphasized by the lack of effect of patient factors including BMI and sex on a success of skull base reconstruction regardless of subsite.

#### Postoperative meningitis and sinusitis

When considering skull base reconstruction and risk of postoperative infections, sinusitis and meningitis are fortunately uncommon complications. In the present study, there was an overall meningitis rate of 2.7% (19/687), which is comparable to the current literature. In a recent systematic review evaluating risk of meningitis following expanded ESBS, the overall incidence of meningitis was found to be 1.8% (80). Unsurprisingly, when postoperative CSF leak was encountered, the risk of meningitis increased compared to those without a postoperative CSF leak (13% vs 0.1%). This emphasizes the importance of successful skull base reconstruction preventing postoperative CSF leak. In the subanalysis comparing ACF and PCF, rates of postoperative meningitis were not statistically different (1.7% vs 1.0%). In the present study, ACF (5.2%) and PCF (4.9%) defects had the highest meningitis rates, but the difference was not significantly different from meningitis arising from sellar (2.3%) and SS (1.7%) surgery. Additionally, we found there were inconsistent factors associated with meningitis, including large tumor size for ACF defects, use of dissolvable nasal packing in SS defects, and highflow intraoperative CSF leaks in PCF defects.

Our study demonstrated an incidence of postoperative sinusitis occurred in 4.4% of cases and was statistically associated with sellar defects when synthetic grafts or nondissolvable packing was used as part of the reconstruction. Importantly, no factors predicted increased risk of sinonasal infection in ACF, SS, or PCF reconstruction. Nondissolvable packing may act as a foreign bodies, with associated risk of local infection. Little et al. examined factors associated with sinonasal quality of life following endoscopic transsphenoidal surgery in 100 patients and found that nasal packing was associated with a higher incidence of postoperative mucopurulence <sup>(81)</sup>. Additionally, mucopurulence resolution in patients with nasal packing occurred significantly later postoperatively (6 weeks) compared to patients without (2 weeks), with presumed impact on patient quality of life. Additionally, Asmaro et al. evaluated 73 consecutive patients after skull base reconstruction without sinonasal packing for CSF leak <sup>(69)</sup>. Infectious sinusitis occurred in 2.7% of patients in the first 3 months postoperatively which is less than the incidence in the present study. Ultimately, there remains a paucity of data exploring the use of these materials in skull base reconstruction and association of sinusitis among other outcomes, thereby warranting further investigation (7,10).

#### Limitations

Although this study leveraged data from four different tertiary rhinology centers, it was still limited by its retrospective nature. Certain pathologies, such as pituitary adenoma, were overrepresented as compared to other rarer pathologies (e.g., chordoma). Moreover, certain factors such as history of previous radiation treatment or comorbid sinusitis were not accounted for in our analysis. Finally, our rates of meningitis and sinus infection were low in our cohort, thereby limiting our statistical analyses of these outcomes. However, notable strengths of this study include a larger sample size across different tertiary skull base programs of wide geographic distribution (U.S. West, Midwest, and Northeast) and diversity in surgical techniques and management principles, which permits for comparison groups, as well as granular data collection with consistent reporting of specific layers of reconstruction, which has not been compiled previously.

# Conclusion

In this multicenter observational study of 706 ESBS patients, we found technical factors, such as the use of rigid reconstruction, dural onlay, and tissue sealants, to be independent predictors of postoperative CSF leak in certain skull base subsites. Perioperative factors, such as the use of synthetic grafts for reconstruction and nondissolvable nasal packing postoperatively, may also incur increased risk for meningitis or sinus infection. Success of skull base reconstruction may largely depend on surgeon adherence to basic principles and necessary elements of skull base reconstruction, and to do so in a meticulous and thoughtful manner.

# Authorship contribution

Data acquisition: AA, BFB, TVN, JCP, KMR, MV, DDC, SHT, JCH, NK, RMS, MK, JEM; Data analysis: AA; Data interpretation: AA, BFB, JED, DJL, JGE, RSK, KMP, ARS, MB, MBW, JNP, NDA, FPKH, ECK; Drafting Article: AA, BFB, TVN, JCP; Final Approval: AA, BFB, TVN, JCP, KMR, MV, DDC, SHT, JCH, NK, RMS, MK, JEM, JED, DJL, JGE, RSK, KMP, ARS, MB, MBW, JNP, NDA, FPKH, ECK

# **Conflict of interest**

There are no relevant conflicts of interest.

#### Funding

This work was supported in part by the National Institute of General Medical Sciences of the National Institutes of Health under award number T32GM008620.

#### **Ethics approval**

This study was conducted in accordance with the Institutional Review Boards at University of California - Irvine, University of California - Los Angeles, University of Pennsylvania, University of Cincinnati.

# Availability of data and materials

Data used in this study is available from the corresponding author (ECK) on reasonable request.

# References

- Wang EW, Zanation AM, Gardner PA, Schwartz TH, Eloy JA, Adappa ND, et al. ICAR: endoscopic skull-base surgery. Int Forum Allergy Rhinol. 2019;9(S3):S145–365.
- Zimmer LA, Theodosopoulos PV. Anterior skull base surgery: open versus endoscopic. Curr Opin Otolaryngol Head Neck Surg. 2009 Apr;17(2):75–8.
- Goshtasbi K, Lehrich BM, Abouzari M, Abiri A, Birkenbeuel J, Lan MY, et al. Endoscopic versus nonendoscopic surgery for resection of pituitary adenomas: a national database study. J Neurosurg. 2020 Mar;134(3):816–24.
- Abiri A, Roman KM, Latif K, Goshtasbi K, Torabi SJ, Lehrich BM, et al. Endoscopic versus Nonendoscopic Surgery for Resection of Craniopharyngiomas. World Neurosurg. 2022 Nov;167:e629–38.
- Fraser S, Gardner PA, Koutourousiou M, Kubik M, Fernandez-Miranda JC, Snyderman CH, et al. Risk factors associated with postoperative cerebrospinal fluid leak after endoscopic endonasal skull base surgery. J Neurosurg. 2018;128(4):1066–71.
- Kim JS, Hong SD. Risk factors for postoperative CSF leakage after endonasal endoscopic skull base surgery: a meta-analysis

and systematic review. Rhinology. 2021 Feb;59(1):10–20.

- Pool C, Abiri A, Kuan EC. Skull Base Reconstruction following Surgical Treatment of Sinonasal Malignancies. J Neurol Surg Rep. 2023 Feb 2;84(1):e17–20.
- Harvey RJ, Parmar P, Sacks R, Zanation AM. Endoscopic skull base reconstruction of large dural defects: a systematic review of published evidence. Laryngoscope. 2012;122(2):452–9.
- Soudry E, Turner JH, Nayak JV, Hwang PH. Endoscopic reconstruction of surgically created skull base defects: a systematic review.

Otolaryngol Neck Surg. 2014;150(5):730-8.

- Abiri A, Abiri P, Goshtasbi K, Lehrich BM, Sahyouni R, Hsu FPK, et al. Endoscopic anterior skull base reconstruction: a meta-analysis and systematic review of graft type. World Neurosurg. 2020 Jul;139:460–70.
- Khan DZ, Ali AMS, Koh CH, Dorward NL, Grieve J, Layard Horsfall H, et al. Skull base repair following endonasal pituitary and skull base tumour resection: a systematic review. Pituitary. 2021 Oct;24(5):698–713.
- Esposito F, Dusick JR, Fatemi N, Kelly DF. Graded repair of cranial base defects and cerebrospinal fluid leaks in transsphenoidal surgery. Oper Neurosurg. 2007;60(2):295– 304.
- Hernán MA, Hernández-Díaz S, Werler MM, Mitchell AA. Causal knowledge as a prerequisite for confounding evaluation: an application to birth defects epidemiology. Am J Epidemiol. 2002 Jan 15;155(2):176–84.
- Yoo W, Mayberry R, Bae S, Singh K, Peter He Q, Lillard Jr JW. A study of effects of multicollinearity in the multivariable analysis. Int J Appl Sci Technol. 2014 Oct;4(5):9–19.
- Soudry E, Turner JH, Nayak JV, Hwang PH. Endoscopic reconstruction of surgically created skull base defects: a systematic review. Otolaryngol Head Neck Surg. 2014 May;150(5):730–8.
- Zanation AM, Carrau RL, Snyderman CH, Germanwala A V., Gardner PA, Prevedello DM, et al. Nasoseptal flap reconstruction of high flow intraoperative cerebral spinal fluid leaks during endoscopic skull base surgery. Am J Rhinol Allergy. 2009;23(5):518– 21.
- 17. Eloy JA, Shukla PA, Choudhry OJ, Singh R, Liu JK. Assessment of frontal lobe sagging after endoscopic endonasal transcribriform resection of anterior skull base tumors: Is rigid structural reconstruction of the cranial base defect necessary? The Laryngoscope. 2012;122(12):2652–7.
- Liu JK, Christiano LD, Patel SK, Tubbs RS, Eloy JA. Surgical nuances for removal of olfactory groove meningiomas using the endoscopic endonasal transcribriform approach. Neurosurg Focus. 2011 May 1;30(5):E3.
- Greenfield JP, Anand VK, Kacker A, Seibert MJ, Singh A, Brown SM, et al. Endoscopic endonasal transethmoidal transcribriform transfovea ethmoidalis approach to the anterior cranial fossa and skull base. Neurosurgery. 2010 May;66(5):883–92.
- Kassam AB, Prevedello DM, Carrau RL, Snyderman CH, Thomas A, Gardner P, et al. Endoscopic endonasal skull base surgery: analysis of complications in the authors' initial 800 patients. J Neurosurg. 2011 Jun;114(6):1544–68.
- 21. Eloy JA, Choudhry OJ, Shukla PA, Kuperan AB, Friedel ME, Liu JK. Nasoseptal flap repair after endoscopic transsellar versus expanded endonasal approaches: is there an increased risk of postoperative cerebrospinal fluid leak? Laryngoscope. 2012

Jun;122(6):1219-25.

- 22. Eloy JA, Patel SK, Shukla PA, Smith ML, Choudhry OJ, Liu JK. Triple-layer reconstruction technique for large cribriform defects after endoscopic endonasal resection of anterior skull base tumors. Int Forum Allergy Rhinol. 2013;3(3):204–11.
- Germani RM, Vivero R, Herzallah IR, Casiano RR. Endoscopic reconstruction of large anterior skull base defects using acellular dermal allograft. Am J Rhinol. 2007;21(5):615–8.
- 24. Hadad G, Bassagasteguy L, Carrau RL, Mataza JC, Kassam A, Snyderman CH, et al. A novel reconstructive technique after endoscopic expanded endonasal approaches: Vascular pedicle nasoseptal flap. Laryngoscope. 2006;116(10):1882–6.
- Esposito F, Dusick JR, Fatemi N, Kelly DF. Graded repair of cranial base defects and cerebrospinal fluid leaks in transsphenoidal surgery. Oper Neurosurg. 2007 Apr;60(4):295.
- 26. Chaskes MB, Fastenberg JH, Vimawala S, Nyquist GF, Rabinowitz MR, Chitguppi C, et al. A simple onlay sellar reconstruction does not increase the risk of postoperative cerebrospinal fluid leak in well-selected patients. J Neurol Surg Part B Skull Base. 2021 Jul;82(Suppl 3):e231–5.
- Sanders-Taylor C, Anaizi A, Kosty J, Zimmer LA, Theodosopoulos PV. Sellar Reconstruction and rates of delayed cerebrospinal fluid leak after endoscopic pituitary surgery. J Neurol Surg Part B Skull Base. 2015 Mar 2;281–5.
- Mehta GU, Oldfield EH. Prevention of intraoperative cerebrospinal fluid leaks by lumbar cerebrospinal fluid drainage during surgery for pituitary macroadenomas: clinical article. J Neurosurg. 2012 Jun 1;116(6):1299–303.
- 29. Romero ADCB, Nora JE, Topczewski TE, Aguiar PHP de, Alobid I, Rodriguéz EF. Cerebrospinal fluid fistula after endoscopic transsphenoidal surgery: experience in a spanish center. Arq Neuropsiquiatr. 2010 Jun;68(3):414–7.
- Nishioka H, Izawa H, Ikeda Y, Namatame H, Fukami S, Haraoka J. Dural suturing for repair of cerebrospinal fluid leak in transnasal transsphenoidal surgery. Acta Neurochir (Wien). 2009 Nov;151(11):1427–30.
- Berker M, Aghayev K, Yücel T, Hazer DB, Onerci M. Management of cerebrospinal fluid leak during endoscopic pituitary surgery. Auris Nasus Larynx. 2013 Aug;40(4):373–8.
- Seda L, Camara RB, Cukiert A, Burattini JA, Mariani PP. Sellar floor reconstruction after transsphenoidal surgery using fibrin glue without grafting or implants: technical note. Surg Neurol. 2006 Jul;66(1):46–9.
- 33. Jalessi M, Sharifi G, Mirfallah Layalestani MR, Amintehran E, Yazdanifard P, Rezaee Mirghaed O, et al. Sellar reconstruction algorithm in endoscopic transsphenoidal pituitary surgery: experience with

240 cases. Med J Islam Repub Iran. 2013 Nov;27(4):186–94.

- 34. Hebert AM, Kuan EC, Wang MB, Snyderman CH, Gardner PA, Bergsneider M, et al. An algorithm for the use of free tissue graft reconstruction in the endoscopic endonasal approach for pituitary tumors. World Neurosurg. 2023 Jul:175:e465-e472.
- Kuan EC, Yoo F, Patel PB, Su BM, Bergsneider M, Wang MB. An algorithm for sellar reconstruction following the endoscopic endonasal approach: a review of 300 consecutive cases. J Neurol Surg Part B Skull Base. 2018 Apr;79(2):177–83.
- Leng LZ, Brown S, Anand VK, Schwartz TH. "Gasket-seal" watertight closure in minimal-access endoscopic cranial base surgery. Neurosurgery. 2008 May;62(5 Suppl 2):ONSE342-343.
- Cavallo LM, Messina A, Esposito F, Divitiis O de, Fabbro MD, Divitiis E de, et al. Skull base reconstruction in the extended endoscopic transsphenoidal approach for suprasellar lesions. J Neurosurg. 2007 Oct 1;107(4):713– 20.
- Garcia-Navarro V, Anand VK, Schwartz TH. Gasket seal closure for extended endonasal endoscopic skull base surgery: efficacy in a large case series. World Neurosurg. 2013 Nov;80(5):563–8.
- Khatiwala RV, Shastri KS, Peris-Celda M, Kenning T, Pinheiro-Neto CD. Endoscopic endonasal reconstruction of high-flow cerebrospinal fluid leak with fascia lata "button" graft and nasoseptal flap: surgical technique and case series. J Neurol Surg Part B Skull Base. 2020 Dec;81(6):645–50.
- 40. Farzal Z, Lemos-Rodriguez AM, Rawal RB, Overton LJ, Sreenath SB, Patel MR, et al. The reverse-flow facial artery buccinator flap for skull base reconstruction: key anatomical and technical considerations. J Neurol Surg Part B Skull Base. 2015 Jun 1;432–9.
- Kim GG, Hang AX, Mitchell C, Zanation AM. Pedicled extranasal flaps in skull base reconstruction. Adv Otorhinolaryngol. 2013;74:71–80.
- 42. Patel MR, Stadler ME, Snyderman CH, Carrau RL, Kassam AB, Germanwala AV, et al. How to choose? endoscopic skull base reconstructive options and limitations. Skull Base. 2010 Nov;20(6):397–404.
- Shah RN, Surowitz JB, Patel MR, Huang BY, Snyderman CH, Carrau RL, et al. Endoscopic pedicled nasoseptal flap reconstruction for pediatric skull base defects. Laryngoscope. 2009 Jun;119(6):1067–75.
- 44. Carrabba G, Dehdashti AR, Gentili F. Surgery for clival lesions: open resection versus the expanded endoscopic endonasal approach. Neurosurg Focus. 2008 Dec 1;25(6):E7.
- Saito K, Toda M, Tomita T, Ogawa K, Yoshida K. Surgical results of an endoscopic endonasal approach for clival chordomas. Acta Neurochir (Wien). 2012 May;154(5):879–86.
- Stippler M, Gardner PA, Snyderman CH, Carrau RL, Prevedello DM, Kassam AB. Endoscopic endonasal approach for

clival chordomas. Neurosurgery. 2009 Feb;64(2):268–77; discussion 277-278.

- Wang EW, Zanation AM, Gardner PA, Schwartz TH, Eloy JA, Adappa ND, et al. ICAR: endoscopic skull-base surgery. Int Forum Allergy Rhinol. 2019;9(S3):S145–365.
- Ahmed OH, Marcus S, Tauber JR, Wang B, Fang Y, Lebowitz RA. Efficacy of perioperative lumbar drainage following endonasal endoscopic cerebrospinal fluid leak repair. Otolaryngol Head Neck Surg Off J Am Acad Otolaryngol-Head Neck Surg. 2017 Jan;156(1):52–60.
- 49. D'Anza B, Tien D, Stokken JK, Recinos PF, Woodard TR, Sindwani R. Role of lumbar drains in contemporary endonasal skull base surgery: meta-analysis and systematic review. Am J Rhinol Allergy. 2016 Nov;30(6):430–5.
- Guo X, Zhu Y, Hong Y. Efficacy and safety of intraoperative lumbar drain in endoscopic skull base tumor resection: a meta-analysis. Front Oncol. 2020 May 7:10:606.
- Mehta GU, Oldfield EH. Prevention of intraoperative cerebrospinal fluid leaks by lumbar cerebrospinal fluid drainage during surgery for pituitary macroadenomas. J Neurosurg. 2012 Jun;116(6):1299–303.
- Birkenbeuel JL, Abiri A, Warner DC, Nguyen E, Marquina S, Gowda S, et al. Lumber drain morbidity in endonasal endoscopic skull base surgery. J Clin Neurosci. 2022 Jul;101:1–8.
- 53. Abiri A, Patel TR, Nguyen E, Birkenbeuel JL, Tajudeen BA, Choby G, et al. Postoperative protocols following endoscopic skull base surgery: an evidence-based review with recommendations. Int Forum Allergy Rhinol. 2023 Jan;13(1):42-71.
- Albu S, Emanuelli E, Trombitas V, Florian IS. Effectiveness of lumbar drains on recurrence rates in endoscopic surgery of cerebrospinal fluid leaks. Am J Rhinol Allergy. 2013;27(6):e190-4.
- Adams AS, Russell PT, Duncavage JA, Chandra RK, Turner JH. Outcomes of endoscopic repair of cerebrospinal fluid rhinorrhea without lumbar drains. Am J Rhinol Allergy. 2016 Nov;30(6):424–9.
- Kitchel SH, Eismont FJ, Green BA. Closed subarachnoid drainage for management of cerebrospinal fluid leakage after an operation on the spine. J Bone Jt Surg - Ser A. 1989;71(7):984–7.
- Scheithauer S, Bürgel U, Bickenbach J, Häfner H, Haase G, Waitschies B, et al. External ventricular and lumbar drainageassociated meningoventriculitis: prospective analysis of time-dependent infection rates and risk factor analysis. Infection. 2010;38(3):205–9.
- 58. Zwagerman NT, Wang EW, Shin SS, Chang YF, Fernandez-Miranda JC, Snyderman CH, et al. Does lumbar drainage reduce postoperative cerebrospinal fluid leak after endoscopic endonasal skull base surgery? A prospective, randomized controlled trial. J Neurosurg. 2018 Oct 1:1-7.

- Hannan CJ, Kewlani B, Browne S, Javadpour M. Multi-layered repair of high-flow CSF fistulae following endoscopic skull base surgery without nasal packing or lumbar drains: technical refinements to optimise outcome. Acta Neurochir (Wien). 2023 Aug;165(8):2299-2307.
- Kitano M, Taneda M. Subdural patch graft technique for watertight closure of large dural defects in extended transsphenoidal surgery. Neurosurgery. 2004 Mar;54(3):653– 61.
- Cappabianca P, Esposito F, Magro F, Cavallo LM, Solari D, Stella L, et al. Natura abhorret a vacuo--use of fibrin glue as a filler and sealant in neurosurgical "dead spaces". Technical note. Acta Neurochir (Wien). 2010 May;152(5):897–904.
- Burkett CJ, Patel S, Tabor MH, Padhya T, Vale FL. Polyethylene glycol (PEG) hydrogel dural sealant and collagen dural graft matrix in transsphenoidal pituitary surgery for prevention of postoperative cerebrospinal fluid leaks. J Clin Neurosci. 2011 Nov;18(11):1513–7.
- Eloy JA, Choudhry OJ, Friedel ME, Kuperan AB, Liu JK. Endoscopic nasoseptal flap repair of skull base defects: is addition of a dural sealant necessary? Otolaryngol Neck Surg. 2012 Jul;147(1):161–6.
- Pereira EAC, Grandidge CA, Nowak VA, Cudlip SA. Cerebrospinal fluid leaks after transsphenoidal surgery – Effect of a polyethylene glycol hydrogel dural sealant. J Clin Neurosci. 2017 Oct;44:6–10.
- 65. Ganesh PB, Basavarajaiah BM, Rudrappa BA, Kasaragod SK. Cerebrospinal fluid rhinorrhoea: does fibrin glue change the surgical outcome? J Laryngol Otol. 2020 Jul;134(7):582–5.
- 66. Laufer I, Anand VK, Schwartz TH. Endoscopic, endonasal extended transsphenoidal, transplanum transtuberculum approach for resection of suprasellar lesions. J Neurosurg. 2007 Mar;106(3):400–6.
- Sautter NB, Batra PS, Citardi MJ. Endoscopic management of sphenoid sinus cerebrospinal fluid leaks. Ann Otol Rhinol Laryngol. 2008 Jan;117(1):32-9.
- Mohindra S, Mohindra S, Gupta K. Endoscopic repair of CSF rhinorrhea: necessity of fibrin glue. Neurol India. 2013;61(4):396.
- Asmaro K, Yoo F, Yassin-Kassab A, Bazydlo M, Robin AM, Rock JP, et al. Sinonasal packing is not a requisite for successful cerebrospinal fluid leak repair. J Neurol Surg Part B Skull Base. 2022 Oct;83(05):476–84.
- Lee TJ, Huang CC, Chuang CC, Huang SF. Transnasal endoscopic repair of cerebrospinal fluid rhinorrhea and skull base defect: ten-year experience: Laryngoscope. 2004 Aug;114(8):1475–81.
- Khan DZ, Marcus HJ, Horsfall HL, Bandyopadhyay S, Schroeder BE, Patel V, et al. CSF rhinorrhoea after endonasal intervention to the skull base (CRANIAL) - part 1: multicenter pilot study. World Neurosurg.

2021 May;149:e1077-89.

- Yano S, Tsuiki H, Kudo M, Kai Y, Morioka M, Takeshima H, et al. Sellar repair with resorbable polyglactin acid sheet and fibrin glue in endoscopic endonasal transsphenoidal surgery. Surg Neurol. 2007 Jan;67(1):59–64.
- Horiguchi K, Murai H, Hasegawa Y, Hanazawa T, Yamakami I, Saeki N. Endoscopic endonasal skull base reconstruction using a nasal septal flap: surgical results and comparison with previous reconstructions. Neurosurg Rev. 2010 Apr;33(2):235–41.
- Bosnjak R, Benedicic M, Vittori A. Early outcome in endoscopic extended endonasal approach for removal of supradiaphragmatic craniopharyngiomas: a case series and a comprehensive review. Radiol Oncol. 2013 Sep 1;47(3):266–79.
- Hsu AK, Singh A, Bury S, Schwartz TH, Anand VK, Kacker A. Endoscopic cerebrospinal fluid leak closure in an infected field. Am J Rhinol Allergy. 2015 Jul;29(4):305–8.
- Zeden JP, Baldauf J, Schroeder HWS. Repair of the sellar floor using bioresorbable polydioxanone foils after endoscopic endonasal pituitary surgery. Neurosurg Focus. 2020 Jun;48(6):E16.
- 77. Tosaka M, Prevedello DM, Yamaguchi R, Fukuhara N, Miyagishima T, Tanaka Y, et al. Single-layer fascia patchwork closure for the extended endoscopic transsphenoidal transtuberculum transplanum approach: deep suturing technique and preliminary results. World Neurosurg. 2021 Nov;155:e271–84.
- Solari D, d'Avella E, Agresta G, Catapano D, D'Ecclesia A, Locatelli D, et al. Endoscopic endonasal approach for infradiaphragmatic craniopharyngiomas: a multicentric Italian study. J Neurosurg. 2022 Jul 1;1–11.
- Yu S, Karsy M, Prashant GN, Barton B, Rosen MR, Parkes W, et al. Minimally invasive endoscopic approaches to pediatric skull base pathologies. Int J Pediatr Otorhinolaryngol. 2022 Nov;162:111332.
- Lai LT, Trooboff S, Morgan MK, Harvey RJ. The risk of meningitis following expanded endoscopic endonasal skull base surgery: a systematic review. J Neurol Surg Part B Skull Base. 2014 Feb;75(1):18–26.
- Little AS, Kelly D, Milligan J, Griffiths C, Prevedello DM, Carrau RL, et al. Predictors of sinonasal quality of life and nasal morbidity after fully endoscopic transsphenoidal surgery. J Neurosurg. 2015 Jun 1;122(6):1458– 65.
- 82. Abiri A, Nguyen TV, Li JY, et al. The impact of nasal packing on skull base reconstruction and quality-of-life outcomes following endoscopic skull base surgery. Int Forum Allergy Rhinol. 2023 Jul 28.
- Phillips KM, Turner MT, Kuan EC. Preoperative, technical, and postoperative considerations for skull base reconstruction: a practical review of critical concepts. Rhinology. 2023;61(5):386-403.

Edward C. Kuan, MD, MBA Department of Otolaryngology Head and Neck Surgery University of California Irvine Medical Center 101 The City Drive South Orange, CA 92868 USA Tel: +1-714 456-5753 Fax: +1-714 456-5747 E-mail: eckuan@uci.edu

Arash Abiri<sup>1</sup>, Benjamin F. Bitner<sup>1</sup>, Theodore V. Nguyen<sup>1</sup>, Jonathan C. Pang<sup>1</sup>, Kelsey M. Roman<sup>1</sup>, Milind Vasudev<sup>1</sup>, Dean D. Chung<sup>1</sup>, Siddhant H. Tripathi<sup>3</sup>, Jacob C. Harris<sup>4</sup>, Nikitha Kosaraju<sup>5</sup>, Ryan M. Shih<sup>6</sup>, Myungjun Ko<sup>6</sup>, Jessa E. Miller<sup>5</sup>, Jennifer E. Douglas<sup>4</sup>, Daniel J. Lee<sup>7</sup>, Jacob G. Eide<sup>8</sup>, Rijul S. Kshirsagar<sup>9</sup>, Katie M. Phillips<sup>3</sup>, Ahmad R. Sedaghat<sup>3</sup>, Marvin Bergsneider<sup>6</sup>, Marilene B. Wang<sup>5</sup>, James N. Palmer<sup>4</sup>, Nithin D. Adappa<sup>4</sup>, Frank P.K. Hsu<sup>2</sup>, Edward C. Kuan<sup>1,2</sup>

<sup>1</sup> Department of Otolaryngology-Head and Neck Surgery, University of California, Irvine, Orange, CA, USA

<sup>3</sup> Department of Otolaryngology-Head and Neck Surgery, University of Cincinnati College of Medicine, Cincinnati, OH, USA

<sup>4</sup> Department of Otorhinolaryngology-Head and Neck Surgery, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

- <sup>5</sup> Department of Head and Neck Surgery, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA
- <sup>6</sup> Department of Neurosurgery, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA
- <sup>7</sup> Department of Otolaryngology-Head and Neck Surgery, Harvard Medical School, Boston, MA, USA
- <sup>8</sup> Department of Otolaryngology-Head and Neck Surgery, Henry Ford Health System, Detroit, MI, USA
- <sup>9</sup> Department of Head and Neck Surgery, Kaiser Permanente Redwood City Medical Center, Redwood City, CA, USA

**Rhinology 62: 3,** 330 - 341, 2024 https://doi.org/10.4193/Rhin23.267

#### \*Received for publication:

July 26, 2023 Accepted: December 28, 2023

#### Assocociate Editor:

Ahmad Sedaghat

This study was presented as a podium at the European Rhinologic Society/International Society of Inflammation and Allergy of the Nose Meeting in Sofia, Bulgaria on June 19, 2023.

<sup>&</sup>lt;sup>2</sup> Department of Neurological Surgery, University of California, Irvine, Orange, CA, USA